

 <p>Agreement on the Conservation of Albatrosses and Petrels</p>	<p><b>Tenth Meeting of the Seabird Bycatch Working Group</b> <i>Virtual meeting, 17 - 19 August 2021 (UTC+10)</i></p> <p><b>Update of incidental capture of seabirds in Argentinean side-haul trawlers</b></p> <p><b><i>Tamini, L. L., Chavez, L. N., Dellacasa, R. F., Crawford R. &amp; E. Frere</i></b></p>
---	--

### SUMMARY

Between April 2008 and July 2015, we conducted a total of 18 trips on five different side-haul trawlers fishing within the Argentine Exclusive Economic Zone, monitoring 486 hauls. We observed 100% of the hauls and monitored trawl cables for 136.7 hours, about 5% of the trawl effort, to identify the levels of seabird bycatch from net entanglements and collisions with trawl cables. A total of 35 net entanglements of White-chinned Petrel *Procellaria aequinoctialis*, Great Shearwater *Ardenna gravis* and Black-browed Albatross *Thalassarche melanophrys* were recorded, all of which occurred during the autumn and winter. Additionally, 656 seabird collisions against trawl cables were recorded including 39 heavy, 96 medium and 521 light. Further, we recorded nine Black-browed Albatross, two Great Shearwater potentially dead. Although in the study fishery the number of deaths in the warp cables could surpass the number of birds incidentally killed in nets, the mortality rate caused by the latter type of interaction far exceeds those observed in nets from other trawl fisheries operating in the Patagonian Shelf. Fortunately, 26% of the seabirds entangled in the net were recovered and released alive, which indicates that awareness and training in safe bird handling and release may improve captured seabird survival rates. The main objectives of this work is to highlight a little-studied source of seabird mortality by entanglement, to generate discussion on potential technical mitigation measures for side-haul trawl fisheries, and to propose crew training in safe handling and release of seabirds as an immediate mitigation measure.

### RESUMEN

Entre abril de 2008 y julio de 2015, realizamos un total de 18 viajes en cinco arrastreros de virado lateral que pescan dentro de la Zona Económica Exclusiva Argentina, monitoreando 486 lances. Observamos el 100% de los lances y monitoreamos los cables de arrastre durante 136,7 horas, aproximadamente el 5% del esfuerzo de arrastre, para identificar los niveles de captura incidental de aves marinas por enredos de redes y colisiones contra los cables de arrastre. Se registraron un total de 35 enredos de Petrel Barba Blanca *Procellaria aequinoctialis*, Pardela Cabeza Negra *Ardenna gravis* y Albatros Ceja Negra *Thalassarche melanophrys*, todos los cuales ocurrieron durante el otoño y el invierno. Además, se registraron 656 colisiones de aves marinas contra cables de arrastre,

incluidas 39 fuertes, 96 intermedias y 521 livianas. Además, registramos nueve Albatros Ceja Negra, dos Pardela Cabeza Negra potencialmente muertos. Si bien, en la pesquería de estudio el número de muertes en los cables de arrastre podría superar el número de aves muertas incidentalmente en las redes, la tasa de mortalidad causada por este último tipo de interacción supera con creces las observadas en las redes de otras pesquerías de arrastre que operan en la Plataforma Patagónica. Afortunadamente, el 26% de las aves marinas enredadas en la red fueron recuperadas y liberadas vivas, lo que indica que la concientización y la capacitación en el manejo y liberación segura de aves pueden mejorar las tasas de supervivencia de las aves capturadas. Los principales objetivos de este trabajo son destacar una fuente poco estudiada de mortalidad de aves marinas por enredos, generar discusión sobre posibles medidas técnicas de mitigación para las pesquerías de arrastre de virado lateral y proponer la capacitación de la tripulación en el manejo seguro y liberación de aves marinas como una medida inmediata de mitigación.

## 1. INTRODUCTION

Seabird bycatch occurs at a rate considered unsustainable for many vulnerable seabird populations, and has a particularly substantial impact on pelagic species (Croxall *et al.* 2012), especially albatrosses and petrels. The cumulative impact of global fisheries on seabird populations became a major conservation concern in the late 1980s (Weimerskirch and Jouventin 1987, Brothers 1991, Murray *et al.* 1993). Although attention focused initially on industrial longlining, bycatch by trawl and artisanal fleets have also been identified as major sources of mortality for many albatrosses and petrels (Sullivan *et al.* 2006b, Favero *et al.* 2011, Croxall *et al.* 2012, Maree *et al.* 2014). Seabird bycatch in trawl fisheries was recorded in the early 1990s (Bartle 1991), but the more cryptic interaction of collisions with trawl cables received less attention than the large numbers of seabirds that were observed hauled aboard, drowned on longline hooks (Weimerskirch *et al.* 2000).

In trawl fisheries, birds are attracted by fish and offal discards, and once the birds are in close proximity to the vessels, mortality occurs due to collisions with trawl cables (metal cables used to tow fishing nets), the “third wire” or netsonde cable (the cable that connects to the net monitor during fishing) or by entanglement in the nets (Weimerskirch *et al.* 2000, González-Zevallos and Yorio 2006, Sullivan *et al.* 2006b). Birds captured in nets or injured/killed after colliding with trawl cables and subsequently hauled onboard are considered to represent the minimum number of those killed during trawling – a large proportion of these birds fall into the water and are not recovered in fishing operations (Weimerskirch *et al.* 2000, Sullivan *et al.* 2006a).

Several factors affect collision rates, including cable characteristics (diameter, material, wear), aerial extent of cables (from the blocks to the entry point in the water), type of discards (i.e. undersized fish, whole heads, guts and offal), quantity and dumping site (i.e. at the stern or side of the vessel) among others (Dietrich and Melvin 2007). Collisions with trawl warp cables has been shown to cause mortality in a large number of seabird species, though long-winged birds, like albatross and *Macronectes* petrels, appear to be the most vulnerable (Sullivan *et al.* 2006b). Entanglements of seabirds occur in trawl fisheries at different operational stages including setting, hauling and trawling (Weimerskirch *et al.* 2000, González-Zevallos and Yorio 2006, Sullivan *et al.* 2006b, Watkins *et al.* 2008). The susceptibility of different seabird groups

to this form of mortality is related to their foraging tactics. Forage-divers (e. g. penguins and cormorants) are more vulnerable to entanglement at depth during trawling and at the start of hauling operations (González-Zevallos *et al.* 2007, Yorio *et al.* 2010), while surface feeders and plunge divers (e. g. petrels and shearwaters) are more vulnerable when the net is close to the surface during setting and hauling (Sullivan *et al.* 2006b, Watkins *et al.* 2008). The *Procellaria* petrels, as well as *Ardenna* shearwaters, are efficient at foraging through pursuit dives and surface-seizing. This feeding method allows them to compete in multi-species feeding assemblages, allowing them to reach fish at depths below those accessible to larger species like albatrosses and *Macronectes* petrels. However, this advantage exposes deeper-diving species to entanglement in fishing gears (Jiménez *et al.* 2012, Jiménez *et al.* 2014, Rollinson *et al.* 2014).

This document, which updates the SBWG8 Doc 10 *Seabirds entanglement in side-haul trawlers: education as a mitigation strategy*, presents a case study of the Argentinean side-haul trawl fishery with the following objectives:

1. outline a detailed description of this poorly-studied haul operation to highlight the key stages that lead to entanglement;
2. describe and analyse the seasonal changes of abundance of highly migratory seabirds that interact with the fishery, to add to existing knowledge of the cumulative impacts these birds face;
3. provide information on the relative importance of seabird mortality through entanglement and collision with trawl cables for this type of fishery;
4. discuss the effectiveness of a short-term strategy for reducing the number of birds bycaught for our study fleet, and potential mitigation measures to be explored.

## 2. MATERIALS AND METHODS

### 2.1. Fishery description and observer coverage

The Patagonian Shelf ecosystem extends from Uruguay to the Isla de los Estados and the Burdwood Bank (35° to 55°S and 50° to 70°W). This ecosystem, a composite area with a unique combination of characteristics, provides fertile conditions that sustain substantial populations of seabirds, seals and whales as well as several coastal and offshore trawl fisheries (Croxall and Wood 2002).

The side-haul trawl fleet of 60 vessels operate year-round between 34°S and 50°S. Vessels lengths range broadly from 20.1 to 59.8 m and can be divided into three potential categories: *small* (20-30 m, 35 vessels), *medium* (30-40 m, 15 vessels) and *large* (40-60 m, 10 vessels), all using demersal trawl nets (120 – 200 mm stretched mesh size at the cod end; 2-3 m and 20-40 m vertical and horizontal aperture respectively). Vessels trawl at a speed of 3.5-4 knots, with trawl durations of between three and four hours, and trips that last a maximum of 18 days. Retained species of commercial interest (target and non-target) are stored whole on ice and no offal is discarded at sea. Usually, discards are composed of small individuals of commercial species, species without current commercial interest and invertebrates (mainly squid, crabs and benthic fauna). In this fleet between April 2008 and July 2015 we conducted a total of 18 cruises on five different vessels from medium (n=3) and large (n=2) categories. Distribution of the trips for this study is similar to the National Onboard Observer Programme pattern, observation in the small strata is difficult by a space and logistic issues (G. Blanco, comm.

pers.). We monitored 486 trawls -observing 100% of the hauls- and achieved 136.7 hours of direct trawl cable observations.

The total annual catch between 2008 and 2015 averaged 112,456 metric tons. This fleet has a dynamic fishing strategy, primarily targeting the Argentine Hake with an average annual catch of 67,323 mt, but also skates (Rajidae, 6,579 mt) and Kingklip (*Genypterus blacodes*, 4,012 mt). Depending on the season, vessels may change fishing gear and/or fishing zone to target Argentine Anchovy (6,954 mt, see Paz *et al.* 2018) and Atlantic Chub Mackerel (*Scomber colias*, 6,248 mt). Some vessels also target Argentine Red Shrimp (*Pleoticus muelleri*).

## 2.2. Seabird assemblage composition and abundance

Seabird abundance to species level was estimated during the trawling operation by one of three experienced seabird observers (LNC, LLT and RFD). Observers conducted approximately 10-min counts from the stern gantry in a 250 m semi-circular sampling area. The overall abundance (total number of individuals in all counts) and abundance per species was calculated. Frequency of occurrence was defined as the percentage of hauls in which each species was observed. The mean number of individuals per haul for the seabirds were compared among seasons with Kruskal-Wallis analysis and the pairwise subsequent comparisons with Dunn's method.

## 2.3. Seabird interactions

We record the entanglement of seabirds in nets during all observed hauls as the net was hauled aboard, noting number of birds caught, species and outcome (dead/unharmed). Owing to the differing likelihoods of recovering bycaught birds between collisions and entanglements, monitoring cable interactions requires a specially designed protocol, which means that estimates of mortality due to collisions are generally poorer than those of mortality due to entanglements (Moore and Żydelis 2008, Watkins *et al.* 2008). The collision of seabirds with trawl cables was recorded using a specially designed protocol during trawling across 251 tows in a total time of 15-90 min per haul ( $\bar{X}$  = 32.1 min, SD = 12.1 min). Data were collected simultaneously for both the starboard and port warps. For collisions with trawl cables the following data were recorded: species, maturity (adult/immature, when this was possible according to the plumage characteristic of the species), severity (heavy/medium/light) and outcome (dead/likely dead/injured/unharmed). The 'likely dead' outcome category was recorded when a bird was seen dragged underwater and not seen again by the observer. The mortality rate associated with trawl cables results from the sum of the dead, likely dead and injured birds. Observations were conducted between daybreak and twilight and throughout the area of fleet operation.

## 3. RESULTS

### 3.1 Seabird assemblage and abundance

Data was gathered from 367 counts over 486 hauls (covering 225 fishing days) in the austral spring (October–December, 70 censuses), winter (July–September, 26), summer (January–March, 169) and autumn (April–June, 102). In total, 23 species of seabird were recorded foraging on discards aft of side-haul trawlers (Table 1). The most frequently observed species (>25% of frequency of occurrence) were the Black-browed Albatross *Thalassarche melanophris* and the White-chinned Petrel *Procellaria aequinoctialis*, followed by the Southern Royal Albatross *Diomedea epomophora*, Great Shearwater *Ardenna gravis* and Southern Giant

Petrel *Macronectes giganteus*. The remainder of the species were observed in low numbers and in <50% of hauls (Table 1).

The abundance of the eight seabird species most frequently associated with this fishery, Black-browed Albatross, White-chinned Petrel, Southern Royal Albatross, Great Shearwater, Southern Giant Petrel, Cape Petrel *Daption capense*, Wilson's Storm Petrel *Oceanites oceanicus*, and Northern Giant Petrel *Macronectes halli*, varied significantly among seasons. Black-browed Albatross and Great Shearwater were more abundant in autumn in comparison with the other seasons. In the same way Southern Royal Albatross were more abundant in summer and winter, Southern Giant Petrel in summer and spring, White-chinned Petrel in summer and autumn and Cape Petrel in spring (pairwise multiple comparison procedures, Dunn's method,  $P < 0.05$ , Table 2).

### **3.2 Seabird interactions**

#### **3.2.1 Net entanglement**

A total of 35 entanglements were recorded, distributed over 15 hauls, including seven hauls with one entanglement, and eight hauls with two to seven entanglements. The entanglement rate of all birds across the 486 observed hauls was 0.071 birds/haul. Four species were recorded entangled including White-chinned Petrel ( $n=20$ ), Great Shearwater ( $n=8$ ), Black-browed Albatross ( $n=6$ ) and a Southern Royal Albatross ( $n=1$ ) at entanglement rates of 0.041; 0.016; 0.012 and 0.002 birds/haul respectively (Table 3). The entanglements were observed exclusively in autumn and winter at a rate of 0.272 and 0.117 birds/haul, respectively. Of the birds recorded entangled, 74% were recovered dead ( $n=26$ , 0.053 birds/haul) while 26% were recovered alive ( $n=9$ , 0.019 birds/haul).

#### **3.2.2. Collision with trawl cables**

From the observed trawls a total of 656 seabird collisions were recorded including 39 heavy, 96 medium and 521 light collisions (Table 4). The majority of collisions were from Black-browed Albatross (418 collisions recorded), Great Shearwater (107), White-chinned Petrel (62), Kelp Gull *Larus dominicanus* (30) and Cape Petrel (21). There were ten or fewer collisions involving the Southern Giant Petrel, Sooty Shearwater *Ardenna grisea*, Southern Royal Albatross, and Southern Fulmar *Fulmarus glacialisoides*. These observations give a total collision rate of 4.8 collisions/hour, calculated as the sum of all interactions (regardless of impact level or outcome) divided by the sum of observation effort. During the observation periods no confirmed mortalities were recorded through collisions; five birds (all Black-browed Albatross) were recorded injured and six birds (four Black-browed Albatross and two Great Shearwater) were recorded as possible fatalities (dragged underwater and not seen again by the observer). A single black browed albatross was recovered dead from trawl cables (Table 3). The mortality rate associated with trawl cables (injured + likely dead) was 0.08 birds/hour (0.002 birds/haul for corpses recovered aboard, Table 3). Collision with trawl cables occurred throughout the year.

## **4. DISCUSSION**

### **4.1 Seabird attendance**

Seabird species in the assemblages attending side-haul trawlers share some features with observations in other studies in the Argentinean EEZ, largely dominated by Procellariiform and Charadriiform species. Fisheries operating close to the coast are attended by a larger numbers of gulls, terns and skuas (Yorio and Caille 1999, González-Zevallos and Yorio 2006, González-

Zevallos *et al.* 2007, Favero *et al.* 2011, Seco Pon *et al.* 2013, Seco Pon 2014, Tamini *et al.* 2015, Paz *et al.* 2018), while those operating in high seas are dominating by albatrosses and petrels. In this study, the Black-browed Albatross was the most frequently observed species, followed by White-chinned Petrel and Great Shearwater. Several studies have identified Black-browed Albatross as the most numerous species associated with Patagonian Shelf fishing vessels, particularly freezer trawlers (Sullivan *et al.* 2006b, González-Zevallos *et al.* 2011, Tamini *et al.* 2015). The presence of Great Shearwater in more than 50% of trawls and in high numbers, especially in autumn (see also González-Zevallos and Yorio 2006, González-Zevallos *et al.* 2011), reaffirms the likelihood that the north Patagonian Shelf is a stop-over in the trans-equatorial migratory route from Tristan da Cunha to the North Atlantic, and/or potentially the result of natural dispersion of some individuals from that archipelago or the small colonies of the Malvinas (Falkland) Islands (Ronconi *et al.* 2018). However, the small sample size in winter may be one of the reasons driving the lack of significant differences in seabird assemblages between seasons. Future studies should attempt to maintain consistency in the number of censuses conducted across seasons to better understand differences in seabird assemblages attending fishing vessels.

#### 4.2 Entanglement details

In trawl fisheries, the hauling operation creates prey availability that otherwise would be naturally inaccessible to shallow divers. While some small albatrosses may dive a few meters, *Procellaria* petrels are proficient (Løkkeborg 2011) and shearwaters are among the best adapted for foot and wing-propelled diving (Burger 2001, Ronconi *et al.* 2010). During hauling operations, seabirds approach the net and mortality through drowning occurs mainly when the wings, legs and/or heads of seabirds become trapped in the mesh. Net interactions associated with the setting and hauling operations occur when the net is floating at the sea surface for extended periods. These periods are longer in this fleet (average 18 minutes, SD = 14, n = 302) than in ramp trawlers, which haul from the stern (average 6 minutes, SD = 1, n = 17), significantly prolonging the availability of the catch to birds (Mann-Whitney U Statistic = 228,500 p <0.001).

Side-haul trawlers are one of the oldest styles of fishing vessel in Argentina's demersal trawl fleet and the hauling and setting operation employed on these vessels is the most widespread across the three segments of the national trawl fleet: wet-fish vessels (called *fresqueros convencionales*), semi-industrial (*costeros*, ~130 vessels) and artisanal bottom trawlers (*rada o ría*, ~380 vessels). The side haul operation involves hauling the cod-end of the net over the side of the vessel, while the trawl warps pass through two blocks suspended at the stern as on other trawlers. Once the trawl doors are retrieved, the vessel performs a tight turn to the starboard side to come alongside the floating net. The haul operation is completed via hauling in two ropes that connect the trawl doors to two different sections of the net, the first around the wings and the second close to the cod end. Each rope gradually brings the cod end closer to the vessel. This system enables manoeuvring and controlling the entire net along the starboard side of the vessel. A boom-rigged mast or crane is then used to hoist the cod-end over the side to empty the catch onto the deck. The final part of the hauling operation is when seabirds are entangled.

#### 4.3 Collisions and entanglements in the Patagonian Shelf

As in other fishing grounds around the world, demersal trawl operations in the Patagonian Shelf fall into two main fleets: ice and freezer trawlers. The seabirds attending vary according

to the type of discards produced, i.e. whole fish on ice trawlers vs. offal plus whole fish on freezers (Weimerskirch *et al.* 2000); observations of seabird trawl cable strike have shown that the presence of offal or whole fish is a key determinant of trawl cable strike rates (Sullivan *et al.* 2006b, Abraham *et al.* 2009). In this study, the observed rate of collision with trawl cables was five times lower than the rate recorded by the entire Argentine ice trawler fleet, including both stern and side-haul vessels (Favero *et al.* 2011), almost seven times lower than recorded in freezer trawlers (Tamini *et al.* 2015), and more than ten times lower than recorded for factory freezer trawlers operating in Malvinas (Falkland) Island waters between August and November when albatross density is high and mean contact rate peaks (55.78 per hour, Sullivan *et al.* 2006a). Our results indicate a rate of 0.29 heavy collisions/hour that is higher when compared to other ice-trawlers (0.04 collision/hour, Favero *et al.* 2011) but insignificant when compared to the 8.36 and 16.8 collisions/hour reported for the freezer trawler fleets in the area (Tamini *et al.* 2015 and Sullivan *et al.* 2006a respectively). For coastal fisheries operating in neighbouring areas, the collision rates are higher for ice trawlers (including side-haul ones), indicating that the distance to the coast or the target specie may be another factors that affects these rates (González-Zevallos *et al.* 2007, Paz *et al.* 2018).

Net entanglement rates have been calculated for a number of fisheries operating on the Patagonian Shelf. Two seasonal coastal fisheries show significant rates of incidental capture of seabirds in the San Jorge Gulf: double-beam trawlers targeting Argentine Red Shrimp (capture rate of 0.048 birds/haul, primarily Magellanic Penguin *Spheniscus magellanicus* and Imperial Cormorant *Phalacrocorax atriceps* (González-Zevallos *et al.* 2011)), and side-haul ice trawlers targeting Argentine Hake (capture rate of 1.2 birds/haul, which also impacts Great Shearwater in addition to the aforementioned penguin and cormorant species, González-Zevallos and Yorio 2006). The high rates in the latter fishery could be due to the combination of seasonality, fishing behaviour and proximity to an area with large seabird colonies (the Marine Protected Area *Parque Interjurisdiccional Marino Costero Patagonia Austral*). High bycatch rates have also been recorded in side-haul vessels targeting Argentine Anchovy (0.55 birds/haul, primarily net mortalities, Paz *et al.* 2018). For side-haul trawlers the rate recorded in this study was 0.071 birds/haul while offshore fisheries have comparatively low levels of net entanglement: 0.009 and 0.013 birds/haul for ice and freezer trawlers respectively (Favero *et al.* 2011, Tamini *et al.* 2015), making it clear that the fishing operation (and the extended time that the net sits on the water's surface) is a key determinant of net entanglement risk, the major bycatch concern in our study fishery. In addition, the species composition of bycaught birds is different between these fleets, with albatrosses primarily affected by ice and freezer trawlers and pursuit-diving species dominating the seabird bycatch composition in our study, particularly Great Shearwater and White-chinned Petrel in autumn.

#### **4.4 Seabird mortality and population impacts**

The three species most impacted by this fishery are captured by other fisheries on the Patagonian shelf, creating the potential for cumulative impacts. Great Shearwater bycatch in demersal longliners is estimated at 0.003 birds/1,000 hooks, a total of 108 birds between 2001-2010 (Favero *et al.* 2013), the offshore Argentine Hake ice trawl fishery of San Jorge Gulf, Argentina, was estimated to kill 2,254 birds in a three month study in 2003 (González-Zevallos and Yorio 2006) and the Argentine Anchovy fishery was observed killed 101 Shearwaters (Great and unidentified) across 172 hauls in 2011-2013 (Paz *et al.* 2018).

Between 2001 and 2007, the bycatch rate of White-chinned Petrel in the Brazilian pelagic longline fleet was estimated as 0.059 birds/1,000 hooks (Bugoni *et al.* 2008). Also, the

Uruguayan fleet caught average of 239 (80-770) birds in the period 2004-2007 and, taking into account the total effort of the pelagic longline fleet, a catch rate of 0.039 birds/1,000 hooks was estimated (Jiménez *et al.* 2010). Further south, in the Argentine demersal longline fleet, the bycatch rate for the period 2001–2010 was 0.012 birds/1,000 hooks, with cumulative annual mortality for the decade estimated at 2,180 ( $\pm$  233) of White-chinned Petrels (Favero *et al.* 2013).

Finally, the fisheries impact to Black-browed Albatross on the Patagonian Shelf is extensive. During the 2001–2007 period, the estimated bycatch rate was 0.126 birds/1000 hooks in Brazilian pelagic longliners (Bugoni *et al.* 2008). In Uruguay, also in pelagic longliners, a mortality of 1,683 (667-3,977) birds was estimated for the 2004-2007 period with a bycatch rate of 0.276 birds/1000 hooks (Jiménez *et al.* 2010). In Argentina, 3,122 ( $\pm$ 336) birds were estimated killed during the 2001–2010 period on demersal longliners, with a bycatch rate of 0.010 birds/1000 hooks (Favero *et al.* 2013). In factory trawlers, an estimated 1,411 birds died annually from cable collisions in the Malvinas/Falkland Island finfish fishery (Sullivan *et al.* 2006b), while for the Argentinean ice-trawler fleet, the annual bycatch rate was 0.012 birds/hour trawling, taking into account net entanglements and cable collisions (Favero *et al.* 2011). A similar bycatch rate (of 0.013 birds/haul) recorded by Tamini *et al.* (2015) for cable collisions alone gave a mortality estimate of 13,548 (8,000-19,673) birds per year (Tamini *et al.* 2015).

Our study supplements this information for these species with a different relevant source of mortality. Based on the number of interactions with nets and cables, we recorded 35 birds entangled across 468 hauls (100% observed) with 20 White-chinned Petrel, eight Great Shearwater and six Black-browed Albatross are included. In addition, we recorded 11 birds (injured and potentially dead plus one recovered bird), composed of nine Black-browed Albatross and two Great Shearwater in 136.7 hours of trawling (based on less than 5% of the trawl time observed). In fact, based on the observed percentage of the trawl effort, the mortality caused by the warp cables could have been substantially higher than the number of birds recorded as potentially dead, and even exceeded the deaths caused by nets. From these data, we do not suspect that many more birds were injured or killed in nets, unlike cables. Considered alone, the bycatch levels of these three species in this fishery are not of population level concern, but should be considered in the context impacts from other fisheries in the Patagonian shelf, some of which are highlighted here. A crucial next step for these three species is improved and harmonized data collection – supported by adequate observer coverage – across fisheries that present a risk, including those outlined at the beginning of this section.

#### **4.5 Mitigation measures**

Physical deterrent mitigation measures – primarily bird-scaring lines – aimed at reducing the number of seabird collisions with warp cable are being used around the Patagonian Shelf with variable levels of implementation (Sullivan *et al.* 2006a, González-Zevallos *et al.* 2007, Snell *et al.* 2012, Tamini *et al.* 2015). Bird-scaring lines could be tested in this fishery, although from a national perspective, priority ought to be given to the freezer trawl fleet because of the substantially higher rate of interactions (Tamini *et al.* 2015).

Solutions that mitigate the impact of trawl fisheries have been identified to varying levels of development (Bull 2007), although no ‘best practice’ measures have been developed to prevent seabirds diving into trawl nets (Løkkeborg 2011). The Agreement on the Conservation of Albatrosses and Petrels Seabird Bycatch Working Group (ACAP SBWG) has identified

“methods to reduce seabirds becoming entangled in nets during hauling” as a priority area for research (ACAP 2016). Measures specifically designed to reduce the incidental capture of seabirds with nets during haul operations were reviewed by Roe (2005). This author suggests the potential use of streamer lines or the reduction of mesh size to minimize entanglements. Streamer lines are impractical in the case of the Argentinean side-haul trawl fishery because trawl speed during hauling is close to zero and the operation is completed over the starboard side, not the stern. The maximum mesh size used by this fleet is larger (200 mm) than that suggested by Roe (2005), so this alternative could be investigated, but with specific consideration of the potential effects on target and non-target fish species. Some alternative mitigation measures, including water deterrent or arrhythmic acoustic sequences to scare seabirds during hauling, may be worthy of testing (Jannot *et al.* 2018). As already noted, seabird entanglements during the hauling operation are a major problem as the net lies slack on the surface for extended periods. Minimizing this time through good operational practice is essential and the rapid retrieval of the net is the key to minimizing seabird interactions and the risk of bycatch. Our observations identified that 26% of the birds entangled were alive and could be released safely. While this only represents a portion of the overall entanglements, it is evident that training crew in seabird handling and release could reduce the number of mortalities. Therefore, in the short term, the development of a training and awareness campaigns to improve crew behaviour provides an opportunity to reduce mortalities due to net entanglements. An additional side benefit of this approach is that it may facilitate engagement with the industry on the development of technical and operational bycatch mitigation approaches.

## REFERENCES

- Abraham, E. R., Pierre, J. P., Middleton, D. A., Cleal, J., Walker, N. A. and Waugh, S. M. (2009). Effectiveness of fish waste management strategies in reducing seabird attendance at a trawl vessel. *Fisheries Research* 95(2-3): 210-219.
- ACAP, S. W. B. G. (2016). Informe del Grupo de Trabajo sobre Captura Secundaria de Aves Marinas CA9 Doc 10 Rev 1. La Serena, Chile, ACAP Agreement of Conservation of Albatrosses and Petrels: 50 p.
- Bartle, J. A. S. (1991). Incidental capture of seabirds in the New Zealand subantarctic squid trawl fishery, 1990. *Bird Conservation International* 1(4): 351-359.
- Brothers, N. (1991). Albatross mortality and associated bait loss in the Japanese longline fishery in the Southern Ocean. *Biological conservation* 55(3): 255-268.
- Bugoni, L., Mancini, P. L., Monteiro, D. S., Nascimento, L. and Neves, T. (2008). Seabird bycatch in the Brazilian pelagic longline fishery and a review of capture rates in the southwestern Atlantic Ocean. *Endangered Species Research* 5: 137-147.
- Bull, L. S. (2007). Reducing seabird bycatch in longline, trawl and gillnet fisheries. *Fish and Fisheries* 8: 31-56.
- Burger, A. E. (2001). Diving depths of shearwaters. *The Auk* 118(3): 755-759.
- Croxall, J. P., Butchart, S. H. M., Lascelles, B., Stattersfield, A. J., Sullivan, B. J., Symes, A. and Taylor, P. (2012). Seabird conservation status, threats and priority actions: a global assessment. *Bird Conservation International* 22(1): 1-34.
- Croxall, J. P. and Wood, A. G. (2002). The importance of the Patagonian Shelf for top predator species breeding at South Georgia. *Aquatic Conservation: Marine and Freshwater Ecosystems* 12(1): 101-118.
- Dietrich, K. S. and Melvin, E. F. (2007). Alaska Trawl Fisheries: Potential Interactions with North Pacific Albatrosses. WSG-TR 07-01. Seattle, WA, Washington Sea Grant: 50 pp.

- Favero, M., Blanco, G., Copello, S., Seco Pon, J. P., Patterlini, C., Mariano-Jelicich, R., García, G. and Berón, M. P. (2013). Seabird bycatch in the Argentinean demersal longline fishery, 2001–2010. *Endangered Species Research* 19(3): 187-199.
- Favero, M., Blanco, G., García, G., Copello, S., Seco Pon, J. P., Frere, E., Quintana, F., Yorio, P., Rabuffetti, F., Cañete, G. and Gandini, P. A. (2011). Seabird mortality associated with ice trawlers in the Patagonian shelf: effect of discards on the occurrence of interactions with fishing gear. *Animal Conservation* 14(2): 131-139.
- González-Zevallos, D. R. and Yorio, P. (2006). Seabird use of discards and incidental captures at the Argentine hake trawl fishery in the Golfo San Jorge, Argentina. *Marine and Ecology Progress Series* 316: 175-183.
- González-Zevallos, D. R., Yorio, P. and Caille, G. (2007). Seabird mortality at trawler warp cables and a proposed mitigation measure: A case of study in Golfo San Jorge, Patagonia, Argentina. *Biological Conservation* 136(1): 108-116.
- González-Zevallos, D. R., Yorio, P. and Svagelj, W. S. (2011). Seabird attendance and incidental mortality at shrimp fisheries in Golfo San Jorge, Argentina. *Marine Ecology Progress Series* 432: 125-135.
- Jannot, J. E., Good, T. P., Tuttle, V. J., Eich, A. M., Fitzgerald, S. M. and (Editors) (2018). US West Coast and Alaska Trawl Fisheries Seabird Cable Strike Mitigation Workshop, November 2017: Summary report. Technical Memorandum NMFS-NWFSC-142. NOAA Department of Commerce: 54 pp.
- Jiménez, S., Abreu, M., Pons, M., Ortiz, M. and Domingo, A. (2010). Assessing the impact of the pelagic longline fishery on albatrosses and petrels in the southwest Atlantic. *Aquatic Living Resources* 23(1): 49-64.
- Jiménez, S., Domingo, A., Abreu, M. and Brazeiro, A. (2012). Bycatch susceptibility in pelagic longline fisheries: are albatrosses affected by the diving behaviour of medium-sized petrels? *Aquatic Conservation: Marine and Freshwater Ecosystems* 22(4): 436-445.
- Jiménez, S., Phillips, R. A., Brazeiro, A., Defeo, O. and Domingo, A. (2014). Bycatch of great albatrosses in pelagic longline fisheries in the southwest Atlantic: Contributing factors and implications for management. *Biological Conservation* 171: 9-20.
- Løkkeborg, S. (2011). Best practices to mitigate seabird bycatch in longline, trawl and gillnet fisheries—efficiency and practical applicability. *Marine and Ecology Progress Series* 435: 285-303.
- Maree, B. A., Wanless, R. M., Fairweather, T. P., Sullivan, B. J. and Yates, O. (2014). Significant reductions in mortality of threatened seabirds in a South African trawl fishery. *Animal Conservation* 17 (6): 520-529.
- Moore, J. E. and Žydelis, R. (2008). Quantifying seabird bycatch: where do we go from here? *Animal Conservation* 11(4): 257-259.
- Murray, T. E., Bartle, J. A. S., Kalish, S. R. and Taylor, P. R. (1993). Incidental capture of seabirds by Japanese southern bluefin tuna longline vessels in New Zealand waters, 1988-1992. *Bird conservation international* 3(3): 181-210.
- Paz, J. A., Seco Pon, J. P., Favero, M., Blanco, G. and Copello, S. (2018). Seabird interactions and by-catch in the anchovy pelagic trawl fishery operating in northern Argentina. *Aquatic Conservation: Marine and Freshwater Ecosystems* 28(4): 850-860.
- Roe, J. (2005). South Georgia Mackerel Icefish (*Champsocephalus gunnari*). Report and Recommendations on Mitigation Use in the Pelagic Icefish Fishery. *Falklands Conservation*: 20 pp.
- Rollinson, D., Dilley, B. and Ryan, P. (2014). Diving behaviour of white-chinned petrels and its relevance for mitigating longline bycatch. *Polar biology* 37(9): 1301-1308.
- Ronconi, R. A., Ryan, P. G. and Ropert-Coudert, Y. (2010). Diving of great shearwaters (*Puffinus gravis*) in cold and warm water regions of the South Atlantic Ocean. *PLoS ONE* 5(11).

- Ronconi, R. A., Schoombie, S., Westgate, A. J., Wong, S. N. P., Koopman, H. N. and Ryan, P. G. (2018). Effects of age, sex, colony and breeding phase on marine space use by Great Shearwaters *Ardenna gravis* in the South Atlantic. *Marine Biology* 165: 58.
- Seco Pon, J. P. (2014). Asociación de aves marinas pelágicas a la flota argentina de arrastre de altura: caracterización integral de las interacciones y desarrollo de una estrategia de conservación para especies con estado de conservación amenazado.
- Seco Pon, J. P., Copello, S., Moretinni, A., Lértora, H. P., Bruno, I., Bastida, J., Mauco, L. and Favero, M. (2013). Seabird and marine-mammal attendance and by-catch in semi-industrial trawl fisheries in near-shore waters of northern Argentina. *Marine and Freshwater Research*. 64: 237-248.
- Snell, K. R. S., Brickle, P. and Wolfaardt, A. C. (2012). Refining Turi lines to further reduce seabird mortality associated with demersal trawlers in the South Atlantic. *Polar biology* 35(5): 677-687.
- Sullivan, B. J., Brickle, P., Reid, T. A., Bone, D. G. and Middleton, D. A. J. (2006a). Mitigation of seabird mortality on factory trawlers: trials of three devices to reduce warp cable strikes. *Polar Biology* 29(9): 745-753.
- Sullivan, B. J., Reid, T. A. and Bugoni, L. (2006b). Seabird mortality on factory trawlers in the Falkland Islands and beyond. *Biological Conservation* 131(4): 495-504.
- Tamini, L. L., Chavez, L. N., Góngora, M. E., Yates, O., Rabuffetti, F. L. and Sullivan, B. J. (2015). Estimating mortality of black-browed albatross (*Thalassarche melanophris*, Temminck, 1828) and other seabirds in the Argentinean factory trawl fleet and the use of bird-scaring lines as a mitigation measure. *Polar Biology* 38(11): 1867-1879.
- Watkins, B. P., Petersen, S. L. and Ryan, P. G. (2008). Interactions between seabirds and deep-water hake trawl gear: an assessment of impacts in South African waters. *Animal Conservation* 11: 247-254.
- Weimerskirch, H., Capdeville, D. and Duhamel, G. (2000). Factors affecting the number and mortality of seabirds attending trawlers and long-liners in the Kerguelen area. *Polar Biology* 23(4): 236-249.
- Weimerskirch, H. and Jouventin, P. (1987). Population Dynamics of the Wandering Albatross, *Diomedea exulans*, of the Crozet Islands: Causes and Consequences of the Population Decline. *Oikos* 49(3): 315-322.
- Yorio, P. and Caille, G. (1999). Seabird Interactions with Coastal Fisheries in Northern Patagonia: Use of Discards and Incidental Captures in Nets *Waterbirds* 22(2): 207-216.
- Yorio, P., Quintana, F., Dell'arciprete, P. and Gonzalez-Zevallos, D. (2010). Spatial overlap between foraging seabirds and trawl fisheries: implications for the effectiveness of a marine protected area at Golfo San Jorge, Argentina. *Bird Conservation International* 20(3): 320-334.

*Noting Article XIII(1)(c) of the Agreement on the Conservation of Albatrosses and Petrels, the references included in the present document are made exclusively for academic/scientific purposes and have no implications whatsoever for recognition of territorial sovereignty or the legal status of a state, territory, area, or their authorities, where relevant.*

## ANNEXES

Table 1: Abundance (total number of individuals in all the census), frequency of occurrence (%) and mean (range in parentheses) per census of seabirds attending side-haul vessels on the Patagonian Shelf during 2008-2015.

<b>Species</b>	<b>Abundance</b>	<b>Frequency of occurrence</b>	<b>Mean (Range)</b>
Black-browed Albatross <i>Thalassarche melanophrys</i>	79,158	99.2	216.3 (0-2,130)
White-chinned Petrel <i>Procellaria aequinoctialis</i>	20,873	96.7	57.0 (0-605)
Southern Royal Albatross <i>Diomedea epomophora</i>	2,813	72.7	7.7 (0-72)
Great Shearwater <i>Ardenna gravis</i>	8,682	53.0	23.7 (0-1536)
Southern Giant Petrel <i>Macronectes giganteus</i>	2,231	51.6	6.1 (0-123)
Cape Petrel <i>Daption capense</i>	2,904	35.8	7.9 (0-390)
Wilson´s Storm Petrel <i>Oceanites oceanicus</i>	826	29.2	2.3 (0-100)
Northern Giant Petrel <i>Macronectes halli</i>	590	25.4	1.6 (0-25)
Kelp Gull <i>Larus dominicanus</i>	1,745	13.4	4.8 (0-310)
Shy/White-capped Albatross <i>Thalassarche cauta/steadii</i>	143	13.4	0.39 (0-20)
Southern Fulmar <i>Fulmarus glacialoides</i>	146	7.9	0.4 (0-63)
Yellow-nosed Albatross <i>Thalassarche chlororhynchos</i>	339	7.1	0.9 (0-155)
Sooty Shearwater <i>Ardenna grisea</i>	61	6.5	2.5 (0-13)
Wandering Albatross <i>Diomedea exulans</i>	35	6.5	0.1 (0-6)
Royal Northern Albatross <i>Diomedea sanfordi</i>	53	6.3	0.15 (0-3)
South American Tern <i>Sterna hirundinacea</i>	284	5.4	0.8 (0-150)
Arctic Jaeger <i>Stercorarius parasiticus</i>	31	4.1	0.09 (0-4)
Brown Skua <i>Catharacta Antarctica</i>	43	3.8	0.12 (0-13)

Chilean Skua <i>Catharacta chilensis</i>	13	3.5	0.04 (0-1)
Grey-headed Albatross <i>Thalassarche chrysostoma</i>	12	2.2	0.03 (0-3)
Slender-billed prion <i>Pachyptila belcheri</i>	4	1.1	0.01 (0-1)
Manx Shearwater <i>Puffinus puffinus</i>	1	0.3	0.003 (0-1)
Spectacled Petrel <i>Procellaria conspicillata</i>	1	0.3	0.003 (0-1)

Table 2: Average abundance (range in parentheses) of individuals per haul for the eight most frequent seabird species attending side-haul trawlers on the Patagonian Shelf during 2008-2015 plus mean total number of birds per haul and the total number of species attending. The seasons in which the first five species were most abundant (Dunn's method) are highlighted with shaded cells. All Kruskal-Wallis tests shown significant differences ( $p < 0.001$ ).

	<b>Summer n=168</b>	<b>Autumn n=102</b>	<b>Winter n=26</b>	<b>Spring n=70</b>	<b>Kruskal-Wallis H</b>
Black-browed Albatross	122.9 (0-1,413)	387.2 (5-2,130)	214.4 (30-950)	192.1 (2-1,470)	32.497
White-chinned Petrel	79.3 (0-605)	48.2 (0-245)	40.7 (2-110)	22.6 (0-180)	45.025
Southern Royal Albatross	11.2 (0-72)	3.7 (0-53)	11.9 (0-50)	3.5 (0-18)	64.194
Great Shearwater	7.6 (0-80)	70.2 (0-1,536)	0	3.5 (0-40)	37.539
Southern Giant Petrel	10.0 (0-64)	0.5 (0-16)	0.6 (0-10)	7.0 (0-123)	102.073
Cape Petrel	0.01 (0-1)	4.0 (0-60)	10.2 (0-40)	31.8 (0-390)	204.169
Wilson's Storm Petrel	2.9 (0-100)	1.0 (0-24)	0.4 (0-2)	3.2 (0-21)	25.224
Northern Giant Petrel	1.7 (0-25)	0.6 (0-13)	0.6 (0-4)	3.3 (0-25)	23.171
Mean abundance	240.7 (2-1,413)	518.6 (15-2,271)	280.0 (44-975)	298.8 (22-1,669)	
Total species	20	18	11	19	

Table 3: Seabirds hauled onboard during 486 hauls (and hauls by season) in side-haul trawler vessels on the Patagonian Shelf during 2008-2015. Birds returned alive in brackets.

<b>Species</b>	<b>Contact with</b>	<b>Summer n=195</b>	<b>Autumn n=92</b>	<b>Winter n=94</b>	<b>Spring n=105</b>	<b>Total n=486</b>	<b>Rates</b>
White-chinned Petrel	Net	-	9	11	-	20 (6)	0.041
	Warp cable	-	-	-	-	-	-
Great Shearwater	Net	-	8	-	-	8	0.016
	Warp cable	-	-	-	-	-	-
Black-browed Albatross	Net	-	6	-	-	6 (3)	0.012
	Warp cable	-	1	-	-	1	0.002
Southern royal Albatross	Net	-	1	-	-	1	0.002
	Warp cable	-	-	-	-	-	-
<b>TOTAL</b>	Net	-	24	11	-	35	<b>0.071</b>
	Warp cable	-	1	-	-	1	<b>0.002</b>

Table 4: Seasonal observation effort displayed as hours, and seabird collisions represented as the numbers of birds and rates (birds/hour) recorded from 486 trawls on industrial side-haul trawl vessels on the Patagonian Shelf between 2008 and 2015.

<b>Seasons</b>	<b>Hours</b>	<b>Heavy</b>		<b>Medium</b>		<b>Light</b>		<b>Total</b>	
		<b>#</b>	<b>Rate</b>	<b>#</b>	<b>Rate</b>	<b>#</b>	<b>Rate</b>	<b>#</b>	<b>Rate</b>
Winter	7.0	0	0.00	3	0.43	20	2.85	23	3.29
Autumn	44.7	18	0.40	31	0.69	315	7.05	364	8.14
Spring	25.5	10	0.39	34	1.33	86	3.37	130	5.09
Summer	59.5	11	0.18	28	0.47	100	1.68	139	2.33
<b>TOTAL</b>	136.7	39	0.29	96	0.70	521	3.81	656	4.80