



Agreement on the Conservation of Albatrosses and Petrels

Fourth Meeting of Seabird Bycatch Working Group

Guayaquil, 22 – 24 August 2011

AC5 Report: Annex 10 — Review of seabird bycatch mitigation measures for demersal longline fishing

Secretariat

‘This paper is presented for consideration by ACAP and may contain unpublished data, analyses, and/or conclusions subject to change. Data in this paper shall not be cited or used for purposes other than the work of the ACAP Secretariat, ACAP Advisory Committee or their subsidiary Working Groups without the permission of the original data holders.’

ANNEX 10: REVIEW OF SEABIRD BYCATCH MITIGATION MEASURES FOR DEMERSAL LONGLINE FISHING

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
1. Avoiding peak areas and periods of seabird foraging activity					
Night setting	(Ashford et al. 1995; Cherel et al. 1996; Moreno et al. 1996; Barnes et al. 1997; Ashford & Croxall 1998; Weimerskirch et al. 2000; Belda & Sánchez 2001; Nel et al. 2002; Ryan & Watkins 2002; Sánchez & Belda 2003; Reid et al. 2004)	Bright moonlight and decklights reduce the effectiveness of this mitigation measure (Cherel et al. 1996; Klaer & Polacheck 1998). Not as effective for crepuscular/nocturnal foragers such as the white-chinned petrel but even for these species night setting is more effective than setting during the day (Ashford et al. 1995; Gómez Laich et al. 2006; Weimerskirch et al. 2000; Nel et al. 2002). In order to maximise effectiveness of this mitigation measure, decklights should be off or kept to an absolute minimum, and used in combination with additional mitigation measures, especially when setting in bright moonlight conditions. Night setting is not a practical option for fisheries operating at high latitudes during summer. Setting should be completed at least 3 hours before sunrise to avoid the predawn activity white-chinned petrels (Barnes et al. 1997)	Recommend combination with bird scaring lines and weighted lines, especially to reduce incidental mortality of birds that forage at night	Effect of night setting on catch rates of target species for different fisheries.	Night defined as the period between the times of nautical twilight (nautical dark to nautical dawn)

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
Area and seasonal closures	<p>A number of studies have reported marked seasonality in seabird bycatch rates, with the majority of deaths taking place during the breeding season (Moreno et al. 1996; Ryan et al. 1997; Ashford & Croxall 1998; Ryan & Purves 1998; Ryan & Watkins 1999; Ryan & Watkins 2000; Weimerskirch et al. 2000; Kock 2001; Nel et al. 2002; Ryan & Watkins 2002; Croxall & Nicol 2004; Reid et al. 2004; Delord et al. 2005). In some studies, mortality has been almost exclusively within the breeding season. Several studies have also shown that proximity to breeding colonies is an important determinant of seabird bycatch rates (Moreno</p>	<p>It is difficult to separate the temporal closure from the increased uptake/ implementation of other mitigation measures, but it is clearly an important and effective management response, especially for high risk areas, and when other measures prove ineffective. There is a risk that temporal/spatial closures could displace fishing effort into neighbouring or other areas which may not be as well regulated, thus leading to increased incidental mortality elsewhere.</p>	<p>Must be combined with other measures, both in the specific areas when the fishing season is opened, and also in adjacent areas to ensure displacement of fishing effort does not lead to a spatial shift in the incidental mortality.</p>	<p>Further information about the seasonal variability in patterns of species abundance, and particularly how these interact with the spatial and temporal characteristics of fishing effort, especially for high risk areas (e.g. adjacent to important breeding colonies). In some studies, incidental mortality has been greatest during the chick-rearing period (Nel et al. 2002; Delord et al. 2005), whereas others have reported highest mortality during the incubation period (Reid et al. 2004). This difference likely relates to where the birds are foraging in relation to fishing effort at the time, and highlights the importance of understanding this interaction. Research is also required to determine the regional impact of closures on catches of</p>	<p>Currently, the area around South Georgia (Islas Georgias del Sur)¹ (CCAMLR Subarea 48.3) is open from May 1st. to Aug. 31st or till established catch limit is reached, as provided for by CCAMLR Conservation Measures in force. (41-02/2007).</p>

¹ “A dispute exists between the Governments of Argentina and the United Kingdom of Great Britain and Northern Ireland concerning sovereignty over the Falkland Islands (Islas Malvinas)”, “South Georgia and the South Sandwich Islands (Islas Georgias del Sur e Islas Sandwich del Sur) and the surrounding maritime areas”.

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	<p>et al. 1996; Nel et al. 2002). The much higher rate of seabird bycatch during the breeding period led to the temporal closure of the fishery in CCAMLR sub-area 48.3 from 1998, which contributed to a ten-fold reduction in seabird bycatch (Croxall & Nicol 2004). Movement of fishing effort away from the Prince Edward Islands coincided with a reduction in seabird bycatch in the sanctioned Prince Edward Island fishery.</p>			<p>target species</p>	
2. Reducing the time baited hooks are near or on the surface and thus available to birds					
<p>Externally weighted lines</p>	<p>(Agnew et al. 2000; Robertson 2000; Melvin et al. 2001; Moreno et al. 2006).</p>	<p>It is important that tension astern is minimised to optimise the sink rate of the line weighting regime. This can be done by preventing hooks snagging on baskets/boxes and by ensuring that weights are released from the vessel before line tension occurs (Robertson et al. 2008a,b). Various methods are used to ensure smooth flow of hooks and avoid entanglements. On autoliners,</p>	<p>Must be combined with other measures, especially bird scaring lines, judicious offal management and/or night setting.</p>	<p>Sink rates and profiles of line weighting regimes may vary according to vessel type, setting speed, how the line is set (relative to the propeller wash for example). It is important that the sink rate relationships of different line weighting regimes are understood for a particular fishery (or fishery method) and that the effectiveness</p>	<p>Global minimum standards not established. Requirements vary by fishery and vessel type. For example, CCAMLR minimum requirements for vessels using the Spanish method of longline fishing are 8.5kg mass at 40m intervals (if rocks are used), 6kg mass at 20m intervals for traditional (concrete) weights, and 5kg weights at 40m intervals for solid steel weights. For autolines, CCAMLR stipulates an</p>

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
		<p>this is achieved by ensuring the correct looping of the line on racks and oiling the line. On the Spanish system it is achieved by correct packing of the lines and hooks and using boxes with smooth edges. Externally attached weights must be attached and removed for each set-haul cycle, which is onerous and potentially hazardous for crew members. Weights made up of rocks enclosed in netting bags and concrete blocks deteriorate and require ongoing maintenance/replacement and monitoring to ensure the required mass is made up (Otley 2005); standard mass weights of steel are better in this respect, both from a handling and compliance perspective (Robertson et al. 2008b). Longlines with externally added weights sink unevenly, faster at the weights than at the midpoint between weights (Robertson 2000). Gear configuration and setting speed influence the sink profiles of the hook lines (Seco Pon et al. 2007), but the principle determinants of sink rates are the mass of the weights and the distance between weights (Robertson et</p>		<p>of the line weighting regime and the sink profile in reducing seabird mortality is tested.</p>	<p>average sink rate of 0.3 m/s to 10 m depth, and a minimum 5kg mass at intervals of no more than 40m. It is also required that weights be released before line tension occurs. In the New Zealand fisheries, a minimum of 4kg (metal weight) or 5kg (non-metal weight) be attached every 60m if the hook bearing line is 3.5mm or greater in diameter, and a minimum of 0.7kg of weight every 60m when the line is less than 3.5mm diameter. The New Zealand minimum standards also include requirements relating to the use of floats.</p>

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
		al. 2008a). See later section on the Chilean longline system.			
Integrated weighting of lines	Apart from the practical advantages of integrated weight (IW) longlines – superior handling qualities and practically inviolable – the IW longlines sink more quickly and uniformly out of reach of most seabirds compared with externally weighted lines. IW longlines have been shown to reduce substantially mortality rates of surface foragers and diving seabirds, while not affecting catch rates of target species (Robertson et al. 2002; Robertson et al. 2003; Robertson et al. 2006; Dietrich et al.	Restricted to autoline vessels. The sink rate of IW longlines can vary depending on vessel type, setting speed and deployment of line relative to propeller wash (Melvin & Wainstein 2006; Dietrich et al. 2008). Setting speed influences the extent of the seabird access window – the area in which most seabirds are still able to access the baited hooks in the absence of bird scaring lines (Dietrich et al. 2008). Use of IW lines is likely to increase the portion of the line on the seafloor, and may lead to increases in the bycatch of vulnerable fish, shark and ray species. This may be mitigated by placing a weight and a float on a 10m line at the point of the dropper line attachment, thus ensuring the line sinks rapidly to 10m, out of reach of vulnerable	Recommended combination with bird scaring lines, judicious offal management and/or night setting.	The relationship between line-weighting regime, setting speed, sink rates/profiles and the seabird access window should be investigated for other fisheries (i.e. those that haven't already been tested –Bering Sea, Alaska, and New Zealand ling fishery) including with additional mitigation measures (particularly bird scaring lines); these investigations would be useful in determining the necessary aerial extent of the bird scaring lines.	Global minimum standards not in place. CCAMLR currently require as a minimum IW lines with a lead core of 50g/m, which is also required in the New Zealand demersal longline fishery. IW must average ≥ 0.24 m/s to 10 m depth. This average rate is less than that for externally weighted non-IW longlines (see above) because IW lines sink with a linear profile from the surface whereas gear with external weights sinks slower between line weights than near the weights.

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	2008)	seabirds, but remains off the seabed (Petersen 2008).			
Side setting	Has not been widely tested in demersal longline fisheries. In trials in the New Zealand ling fishery, side setting appeared to reduce seabird bycatch; however, the results were not convincing and there were practical/operational difficulties, with the line becoming entangled in the propeller (Bull 2007). Sullivan (2004) reported that side setting has been used in some demersal fisheries (e.g. shark fisheries) which have experienced negligible incidental mortality.	Practical difficulties, especially in difficult weather/sea conditions. In many cases it may be difficult and expensive converting the vessel's deck design to employ a side setting system.	In pelagic longline fisheries must be used in combination with other mitigation measures, especially the use of a bird curtain (Gilman et al. 2007), and bird scaring lines.	Largely untested in the demersal fisheries, especially in the Southern Ocean, where the seabird assemblages include proficient diving seabirds. Research urgently needed.	None. Insufficient evidence to recommend this measure

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
Underwater setting funnel/chute	An underwater setting funnel has been tested in demersal longline fisheries in Alaska, Norway and South Africa, with all studies showing a reduction in the mortality rate, although the extent of the reduction varied between studies (Løkkeborg 1998, 2001; Melvin et al. 2001; Ryan & Watkins 2002).	Present design is mainly for a single line system. Results from studies to date have been inconsistent, likely due to the depth at which the device delivers the baited hooks and the diving ability of the seabirds in the fishing area studied. The pitch angles of the vessel, which are influenced by the loading of weight and sea conditions, affect the performance of the funnel (Løkkeborg 2001).	Must be used in conjunction with other mitigation measures – bird scaring lines, weighted lines, night setting and judicious offal management.	Need to investigate improvements to the current design to increase the depth at which the line is set, especially during rough seas. Should also be tested with integrated weight lines to determine whether this improves bycatch reduction. Also need to investigate optimal use of device together with other mitigation measures (bird scaring lines and weighted lines).	Not yet established
Line setter/shooter	Less used in demersal long-line fisheries; variation in the precise method of operation is cause of variation in efficacy. Reduced bycatch of northern fulmars relative to sets with no mitigation measures in trials conducted in Norway, but not significantly (Løkkeborg & Robertson 2002; Løkkeborg 2003). However, seabird bycatch in Alaska increased when a line shooter was used	A significant reduction in seabird bycatch when setting with a line shooter has not been demonstrated. Robertson et al. (2008c) found no significant difference between the sink rates of integrated weight longlines of autoline vessels that were set with and without a line setter in the Ross Sea, and were doubtful that the use of line setters would lead to substantial reductions in interactions between seabirds and longlines. At this stage it should not be seen as a measure.	Currently not considered a mitigation measure.	.	None. Insufficient evidence to recommend this measure

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	(Melvin et al. 2001).				
Thawing bait	See comments for pelagic longline fisheries. In demersal longline fisheries operational considerations ensure baits are thawed to the point where thaw status does not affect sink rates.	No an effective mitigation measure.		There is some evidence that the number of seabirds caught varies according to the species of bait used (Weimerskirch et al. 2000). This should be investigated further.	None. Insufficient evidence to recommend this measure
3. Actively deterring birds from baited hooks					
Single bird scaring line	The use of a single bird scaring line has been shown to be an effective mitigation measure in a range of demersal longline fisheries, especially when used properly (Moreno et al. 1996;	Effective only when streamers are positioned over sinking hooks. Single bird scaring lines can be less effective in strong crosswinds (Løkkeborg 1998; Agnew et al. 2000; Melvin et al. 2001; Melvin et al. 2004). In the event of strong crosswinds, bird scaring lines should be	Effectiveness is increased when used in combination with other measures – e.g. night setting, appropriate weighting of line and judicious offal	The use and specifications/performance standards are fairly well established in demersal longline fisheries. However, there is scope to improve further the effectiveness and practical use of bird scaring lines on	Current minimum standards vary. CCAMLR was the first conservation body that required all longline vessels in its area of application to use bird scaring lines (Conservation Measure 29/X adopted in 1991). The bird scaring line has gone on to become the most commonly

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	<p>Løkkeborg 1998, 2001; Melvin et al. 2001; Smith 2001; Løkkeborg & Robertson 2002; Løkkeborg 2003; Robertson et al., 2008a)</p>	<p>deployed from the windward side. This problem can also be overcome by using paired bird scaring lines (see below).The effectiveness of the bird scaring lines is also dependent on the design, the aerial coverage of the bird scaring line, seabird species present during line setting (proficient divers being more difficult to deter from baits than surface feeding birds) and the proper use of the bird scaring line. The aerial coverage and the position of the bird scaring line relative to the sinking hooks are the most important factors influencing their performance. There have been a few incidents of birds becoming entangled in bird scaring lines (Otley et al. 2007). However it must be stressed that the numbers are minuscule, especially when compared with the number of mortalities recorded in the absence of bird scaring lines. Bird scaring lines remain a highly effective mitigation measure, and efforts should be directed to improving further their design and use so that their effectiveness can be improved further.</p>	<p>management.</p>	<p>individual vessels or vessel type.</p>	<p>applied mitigation measure in longline fisheries worldwide (Melvin et al. 2004). CCAMLR currently prescribes a range of specifications relating to the design and use of bird scaring lines. These include the minimum length of the line (150m), the height of the attachment point on the vessel (7m above the water), and details about streamer lengths and intervals between streamers. Other fisheries have adapted these measures. Some, such as those in New Zealand and Alaska have set explicit standards for the aerial coverage of the bird scaring lines, which varies according to the size of the vessel.</p>

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
Paired or multiple bird scaring lines	Several studies have shown that the use of two or more streamer lines is more effective at deterring birds from baited hooks than a single streamer line (Melvin et al. 2001; Sullivan & Reid 2002; Melvin 2003; Melvin et al. 2004; Reid et al. 2004). The combination of paired streamer lines and IW longlines is considered the most effective mitigation measure in demersal longline fisheries using autoline systems (Dietrich et al. 2008).	Potentially increased likelihood of entanglement with other gear. Use of an effective towed device that keeps lines from crossing surface gear essential to improve adoption and compliance. See also above comment about bird entanglements in bird scaring lines. Manually attached and operated paired or multiple bird scaring lines requires some effort to operate (a 150m double line takes several men to retrieve). One way of overcoming this is to make use of electronic winches.	Effectiveness is increased when used in combination with other measures – e.g. night setting, appropriate weighting of line and judicious offal management.	Further trialling in fisheries which currently only use single streamer lines.	Paired streamer lines required in Alaskan fisheries and encouraged/recommended by CCAMLR, except in the French exclusive economic zone (CCAMLR Subarea 58.6 and Division 58.5.1), where paired streamer lines have been compulsory since 2005. Paired streamer lines have also been required in the Australian longline fisheries off Heard Island since 2003 (Dietrich et al. 2008)
Haul mitigation	The use of a bird exclusion device (BED) such as a Brickle curtain can effectively reduce the incidence of birds becoming foul hooked when the line is being hauled (Brothers et al. 1999; Sullivan 2004; Otley et al. 2007; Reid et al. submitted, Snell et al. in prep.).	Some species, such as the black-browed albatross and cape petrels, can become habituated to the curtain, so it is important to use it strategically – when there are high densities of birds around the hauling bay (Sullivan 2004).	None known		A BED discourages birds from accessing baits during hauling operations and is required in high risk CCAMLR areas. CCAMLR has not specified the exact design, but a BED must: 1) deter birds from flying into the area where the line is being hauled, and 2) prevents birds that are sitting on the surface from swimming into the hauling bay area. Also required in the Falkland Islands (Islas Malvinas) ¹

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
					longline fishery, where the Brickle Curtain is recommended (Snell et al, in prep).
Olfactory deterrents	Dripping shark liver oil on the sea surface behind vessels has been shown to effectively reduce the number of seabirds (restricted to burrow-nesting birds) attending vessels and diving for bait in New Zealand (Pierre & Norden 2006; Norden & Pierre 2007).	The shark liver oil did not deter albatrosses, giant petrels, or Cape Petrels from boats (Norden & Pierre 2007). The potential impact of releasing large amounts of concentrated fish oil into the marine environment is unknown, as is the potential for contaminating seabirds attending vessels and the potential of seabirds to become habituated to the deterrent (Pierre & Norden 2006).	Must be used in combination with other mitigation measures – bird scaring lines at setting, line weighting, night setting and judicious offal management – especially until further testing has been conducted.	Testing should be extended to candidate/suitable species of conservation concern, such as white-chinned petrels and sooty shearwaters. Research is also required to identify the key ingredients in the shark oil that are responsible for deterring seabirds, and the mechanism by which the birds are deterred. The potential “pollution” effects also need to be investigated.	None yet.
4. Reducing attractiveness and visibility of baited hooks and attractiveness of vessel to birds					
Strategic management of offal discharge	Some studies have shown that dumping homogenised offal (which is generally more easily available and thus attractive to seabirds than bait) during setting attracts birds away from the baited line to the side	Although strategic offal discharge has been shown to be effective at reducing seabird bycatch around Kerguelen Island, there are many risks associated with the practice. Offal discharge needs to be continued throughout the setting operation so as to ensure the birds do not move on to the	Not recommended as a mitigation measure at this time. See comments in “Minimum standards”, which are considered best practice approaches to offal	Further information needed on opportunities to manage offal more effectively – considering both practical aspects and seabird bycatch mitigation – in the short and long term.	In CCAMLR demersal fisheries, discharge of offal is prohibited during line setting. During line hauling, storage of waste is encouraged, and if discharged must be discharged on the opposite side of the vessel to the hauling bay. A system to remove fish hooks from offal and fish heads prior to discharge is

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	<p>of the vessel where the offal is being discharged, and thus reduces bycatch of seabirds on the baited hooks (Cherel et al. 1996; Weimerskirch et al. 2000).</p>	<p>baited hooks. This will only be possible in fisheries where line setting is short, and there is sufficient offal to sustain the line-setting period. This measure also has the potential to foul hook birds if offal is discharged with hooks. It is crucial, then, that all offal is checked for hooks before being discharged. Given these risks, and the fact that the presence of offal is a critical factor affecting seabird numbers attending vessels, most fisheries management agencies require that no offal can be discharged during line setting, and that if discarding is necessary at other times it should take place on the side of the vessel opposite to where the lines are being hauled.</p>	<p>management.</p>		<p>required. Similar requirements are prescribed by other demersal longline fisheries (e.g. Falkland Islands (Islas Malvinas)¹, South Africa and New Zealand)</p>
<p>Blue-dyed bait</p>	<p>The performance of this measure has only been tested in the pelagic longline fishery (Boggs 2001; Minami & Kiyota 2004; Gilman et al. 2007; Cocking et al. 2008), and with mixed success.</p>	<p>This measure is only effective with squid bait (Cocking et al. 2008). It has not been tested in demersal fisheries, possibly due to the larger number of hooks deployed and thus the need for considerably more bait (Bull 2007). There is no commercially available dye. Onboard dyeing is practically onerous, especially in inclement weather. In the long-term birds may become</p>	<p>Must be used in combination with other mitigation measures – bird scaring lines, line weighting, night setting and judicious offal management</p>	<p>Need for tests of efficacy and practical feasibility in demersal longline fisheries, especially in the Southern Ocean to determine its effectiveness as a long-term mitigation measure. Research would also need to determine the effect of dyed bait on catches of target species.</p>	<p>Mix to standardized colour placard or specify (e.g. use 'Brilliant Blue' food dye (Colour Index 42090, also known as food additive number E133) mixed at 0.5% for a minimum of 20 minutes).</p>

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
		habituated to blue-dyed bait.			
5. Other					
Hook size and shape	Hook size was found to be an important determinant in seabird bycatch rates of Argentinean and Chilean longline vessels fishing in Subarea 48.3 in the 1995 season, with smaller hooks killing significantly more seabirds than larger hooks (Moreno et al. 1996)	Other than the finding in Moreno et al (1996), little or no work has been conducted to investigate the impact of hood design and shape on seabird bycatch levels.	Unknown, due to insufficient understanding about effectiveness as seabird deterrent.	Further studies required to determine impact on seabird bycatch and on catch of target species	None. Insufficient evidence to recommend this measure.
Gear configuration – Chilean method (linked with sink rates)	A new method of demersal longline fishing, called the Chilean longline method, developed from the Chilean artisanal toothfish fishery, has been shown to reduce significantly seabird bycatch as a consequence of rapid	This is a new system and should be monitored and possibly refined further. Concern has been raised about the excessive discard of unwanted hooks that may be associated with this longline system and the ingestion of these hooks – and discarded fish bearing them - by seabirds (Phillips et al. 2010). The solution to this problem is to ban	One of the few techniques that is effective on its own. Preferably use in combination with bird scaring lines.	Test broader applicability to other fisheries, including effects on fish bycatch. In other fisheries the relationship between weight mass, weight type and sink rate (see Robertson et al., 2008a for the Patagonian toothfish fishery) should be investigated to determine the minimum weight	No global standards yet

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	<p>sink rates compared with traditional longline systems (Moreno et al. 2006; Moreno et al. 2008; Robertson et al. 2008b). This system makes use of net sleeves or ‘cachaloteras’ which slide down over the hooks and captured fish during hauling and thus protect fish from toothed whales. The advantage of this configuration is that hook-bearing lines sink vertically (not horizontally) in the water column in the depths assessable to seabirds, and that hooks are located close to heavy line weights (ensures rapid sinking). This system was first tested on large longline vessels in 2005. Because of the effectiveness of the Chilean longline system in reducing impacts of toothed whales, it is currently</p>	<p>the discarding of hooks as part of the licence conditions, as is already done in many fisheries, and educating fishers, observers and operators to facilitate compliance with such a ban. CCAMLR has produced a weather-proof educational poster on the importance of retaining hooks onboard and distributed the poster to all vessels operating in Convention Area waters.</p>		<p>requirement. The Chilean system is used primarily to prevent depredation of caught fish by cetaceans, the by-product of which is significantly reduced seabird bycatch. Given the possibility that cetaceans may become habituated to the net sleeves over time, it is important that the efficacy of this system at deterring cetaceans continues to be monitored.</p>	

Mitigation measure	Scientific evidence for effectiveness in demersal fisheries	Caveats /Notes	Need for combination	Research needs	Minimum standards
	used in many longline fleets operating in South American waters (Moreno et al. 2008), as well as in the south west Atlantic.				

REFERENCES

- Agnew, D.J., Black, A.D., Croxall, J.P. and Parkes, G.B. 2000. Experimental evaluation of the effectiveness of weighting regimes in reducing seabird by-catch in the longline toothfish fishery around South Georgia. *CCAMLR Science*, 7: 119–131.
- Ashford, J.R., and Croxall, J.P. 1998. An assessment of CCAMLR measures employed to mitigate seabird mortality in longline operations for *Dissostichus eleginoides* around South Georgia. *CCAMLR Science*, 5: 217–230.
- Ashford, J.R., Croxall, J.P., Rubilar, J.S. and Moreno, C.A. 1995. Seabird interactions with longlining operations for *Dissostichus eleginoides* around South Georgia, April to May 1994. *CCAMLR Science*, 2: 111–121.
- Barnes, K.N., Ryan, P.G. and Boix-Hinzen, C. 1997. The impact of the hake *Merluccius spp.* longline fishery off South Africa on procellariiform seabirds. *Biological Conservation*, 82: 227–234.
- Belda, E.J., and Sánchez, A. 2001. Seabird mortality on longline fisheries in the western Mediterranean: factors affecting bycatch and proposed mitigating measures. *Biological Conservation*, 98: 357–363.
- Boggs, C.H. 2001. Deterring albatrosses from contacting baits during swordfish longline sets. Pages 79–94 in E.F. Melvin, and J.K. Parrish, editors. *Seabird Bycatch: Trends, Roadblocks and Solutions*. University of Alaska Sea Grant, AK-SG-01, Fairbanks, AK.
- Brothers, N.P., Cooper, J., and Lokkeborg, S. 1999. The incidental catch of seabirds by longline fisheries: worldwide review and technical guidelines for mitigation. *FAO Fisheries Circular* 937.
- Bull, L.S. 2007. Reducing seabird bycatch in longline, trawl and gillnet fisheries. *Fish and Fisheries*, 8: 31–56.
- Cherel, Y., Weimerskirch, H. and Duhamel, G. 1996. Interactions between longline vessels and seabirds in Kerguelen waters and a method to reduce seabird mortality. *Biological Conservation*, 75: 63–70.
- Cocking, L.J., Double, M.C., Milburn, P.J. and Brando, V.E. 2008. Seabird bycatch mitigation and blue-dyed bait: A spectral and experimental assessment. *Biological Conservation*, 141: 1354–1364.
- Croxall, J.P., and Nicol, S. 2004. Management of Southern Ocean fisheries: global forces and future sustainability. *Antarctic Science*, 16: 569–584.
- Delord, K., Gasco, N., Weimerskirch, H., Barbraud, C. and Micol, T. 2005. Seabird mortality in the Patagonian toothfish longline fishery around Crozet and Kerguelen Islands, 2001-2003. *CCAMLR Science*, 12: 53–80.
- Dietrich, K.S., Melvin, E.F. and Conquest, L. 2008. Integrated weight longlines with paired streamer lines - best practice to prevent seabird bycatch in demersal longline fisheries. *Biological Conservation*, 141: 1793–1805.
- Gilman, E., Brothers, N. and Kobayashi, D.R.. 2007. Comparison of three seabird bycatch avoidance methods in Hawaii-based pelagic longline fisheries. *Fisheries Science*, 73: 208–210.
- Gilman, E., Brothers, N. and Kobayashi, R. 2005. Principles and approaches to abate seabird by-catch in longline fisheries. *Fish and Fisheries*, 6: 35–49.
- Gómez Laich A, Favero, M., Mariano-Jelicich, R., Blanco, G., Cañete, G., Arias, A., Silva Rodriguez, M.P. and Brachetta, H. 2006. Environmental and operational variability affecting the mortality of black-browed albatrosses associated to long-liners in Argentina. *Emu*, 106: 21–28.

- Klaer, N., and Polacheck, T. 1998. The influence of environmental factors and mitigation measures on bycatch rates of seabirds by Japanese longline vessels in the Australian region. *Emu*, 98: 305–306.
- Kock, K.-H. 2001. The direct influence of fishing and fishery-related activities on non-target species in the Southern Ocean with particular emphasis on longline fishing and its impact on albatrosses and petrels - a review. *Reviews in Fish Biology and Fisheries*, 11: 31–56.
- Løkkeborg, S. 1998. Seabird by-catch and bait loss in long-lining using different setting methods. *ICES Journal of Marine Science*, 55: 145–149.
- Løkkeborg, S. 2001. Reducing seabird bycatch in longline fisheries by means of bird-scaring and underwater setting. Pages 33-41 in E.F. Melvin, and J.K. Parrish, editors. *Seabird Bycatch: Trends, Roadblocks and Solutions*. University of Alaska Sea Grant, Fairbanks, AK.
- Løkkeborg, S. 2003. Review and evaluation of three mitigation measures - bird-scaring line, underwater setting and line shooter - to reduce seabird bycatch in the north Atlantic longline fishery. *Fisheries Research*, 60: 11–16.
- Løkkeborg, S., and Robertson, G. 2002. Seabird and longline interactions: effects of a bird-scaring streamer line and line shooter on the incidental capture of northern fulmars *Fulmarus glacialis*. *Biological Conservation*, 106: 359–364.
- Melvin, E.F. 2003. Streamer lines to reduce seabird bycatch in longline fisheries. Washington Sea Grant Program WSG-AS 00-33.
- Melvin, E.F., and Parrish, J.K. editors. 2001. *Seabird bycatch: trends, roadblocks and solutions*. University of Alaska Sea Grant, AK-SG-01-01, Fairbanks, AK.
- Melvin, E.F., Parrish, J.K., Dietrich, K.S. and Hamel, O.S. 2001. Solutions to seabird bycatch in Alaska's demersal longline fisheries. Washington Sea Grant Program. Project A/FP-7. WSG-AS 01-01. University of Washington, Seattle WA.
- Melvin, E.F., and Robertson, G. 2001. Seabird mitigation research in long-line fisheries: Status and priorities for future research and actions. *Marine Ornithology*, 28: 178–181.
- Melvin, E.F., Sullivan, B., Robertson, G. and Wienecke, B. 2004. A review of the effectiveness of streamer lines as a seabird by-catch mitigation technique in longline fisheries and CCAMLR streamer line requirements. *CCAMLR Science*, 11: 189–201.
- Melvin, E.F., and Wainstein, M.D. 2006. Seabird avoidance measures for small Alaskan longline vessels. Project A/FP-7. Washington Sea Grant Program.
- Minami, H., and Kiyota, M. 2004. Effect of blue-dyed bait and tori-pole streamer on reduction of incidental take of seabirds in the Japanese southern bluefin tuna longline fisheries. CCSBT-ERS/0402/08. CCSBT, Canberra.
- Moreno, C.A., Arata, J.A., Rubilar, P., Hucke-Gaete, R. and Robertson, G. 2006. Artisanal longline fisheries in Southern Chile: Lessons to be learned to avoid incidental seabird mortality. *Biological Conservation*. 127: 27–37.
- Moreno C.A., Castro, R., Mujica L.J. and Reyes, P. 2008. Significant conservation benefits obtained from the use of a new fishing gear in the Chilean Patagonian toothfish fishery. *CCAMLR Science*, 15: 79–91.
- Moreno, C.A., Rubilar, P.S. Marschoff, E. and Benzaquen, L. 1996. Factors affecting the incidental mortality of seabirds in the *Dissostichus eleginoides* fishery in the south-west Atlantic (Subarea 48.3, 1995 season). *CCAMLR Science*, 3: 79–91.
- Nel, D.C., Ryan, P.G. and Watkins, B.P. 2002. Seabird mortality in the Patagonian toothfish longline fishery around the Prince Edward Islands, 1996-2000. *Antarctic Science*, 14: 151–161.

- Norden, W.S., and Pierre, J.P. 2007. Exploiting sensory ecology to reduce seabird by-catch. *Emu*, 107: 38–43.
- Otley, H. 2005. Seabird mortality associated with Patagonian toothfish longliners in Falkland Island waters during 2002/03 & 2003/04. Falkland Islands Fisheries Department, Stanley, Falkland Islands.
- Otley, H.M., Reid, T.A. and Pompert, J. 2007. Trends in seabird and Patagonian toothfish *Dissostichus eleginoides* longliner interactions in Falkland Island waters, 2002/03 and 2003/04. *Marine Ornithology*, 35: 47–55.
- Petersen, S.L. 2008. Understanding and mitigating vulnerable bycatch in longline and trawl fisheries off southern Africa. Unpublished PhD thesis, University of Cape Town, Cape Town, South Africa.
- Phillips, R.A, Ridley, C., Reid, K., Pugh, P.J.A., Tuck, G.N. and Harrison, N. 2010. Ingestion of fishing gear and entanglements of seabirds: monitoring and implications for management. *Biological Conservation*, 143: 501–512.
- Pierre, J.P., and Norden, W.S. 2006. Reducing seabird bycatch in longline fisheries using a natural olfactory deterrent. *Biological Conservation*, 130: 406–415.
- Reid, E., Sullivan B., and Clark, J. submitted. Mitigation of seabird captures during hauling in CCAMLR longline fisheries. CCAMLR Science.
- Reid, T.A., Sullivan, B.J., Pompert, J., Enticott, J.W. and Black, A.D. 2004. Seabird mortality associated with Patagonian toothfish (*Dissostichus eleginoides*) longliners in Falkland Islands waters. *Emu*, 104: 317–325.
- Robertson, G., McNeill, M., King, B. and Kristensen, R. 2002. Demersal longlines with integrated weight: a preliminary assessment of sink rates, fish catch success and operational effects. CCAMLR-WG-FSA-02/22. CCAMLR, Hobart.
- Robertson, G., McNeill, M., Smith, N., Wienecke, B., Candy, S. and Olivier, F. 2006. Fast sinking (integrated weight) longlines reduce mortality of white-chinned petrels (*Procellaria aequinoctialis*) and sooty shearwaters (*Puffinus griseus*) in demersal longline fisheries. *Biological Conservation*, 132: 458–471.
- Robertson, G., Moe, E., Haugen, R. and Wienecke, B. 2003. How fast do demersal longlines sink? *Fisheries Research*, 62: 385–388.
- Robertson, G., Moreno, C.A., Crujeiras, J., Wienecke, B., Gandini, P.A., McPherson, G. and Seco Pon, J.P. 2008a. An experimental assessment of factors affecting the sink rates of Spanish-rig longlines to minimize impacts on seabirds. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 17: S102–S121.
- Robertson, G., Moreno, C.A., Gutiérrez, E., Candy, S.G., Melvin, E.G. and Seco Pon, J.P. 2008b. Line weights of constant mass (and sink rates) for Spanish-rig Patagonian toothfish longline vessels. CCAMLR Science, 15: 93–106.
- Robertson, G., Williamson, J., McNeill, M., Candy, S.G. and Smith, N. 2008c. Autoliners and seabird by-catch: do line setters increase the sink rate of integrated weight longlines? CCAMLR Science, 15: 107–114.
- Robertson, G.G. 2000. Effect of line sink rate on albatross mortality in the Patagonian toothfish longline mortality. CCAMLR Science, 7: 133–150.
- Ryan, P. and Watkins, B. 2000. Seabird by-catch in the Patagonian toothfish longline fishery at the Prince Edward Islands: 1999 - 2000. CCAMLR-WG-FSA 00/30. CCAMLR, Hobart.
- Ryan, P.G., Boix-Hinzen, C., Enticott, J.W., Nel, D.C., Wanless, R. and Purves, M. 1997. Seabird mortality in the longline fishery for Patagonian Toothfish at the Prince Edward Islands: 1996 - 1997. CCAMLR-WG-FSA 97/51. CCAMLR, Hobart.

- Ryan, P.G. and Purves, M. 1998. Seabird bycatch in the Patagonian toothfish fishery at Prince Edward Islands: 1997-1998. CCAMLR-WG-FSA 98/36. CCAMLR, Hobart.
- Ryan, P.G. and Watkins, B.P. 1999. Seabird by-catch in the Patagonian toothfish longline fishery at the Prince Edward Islands: 1998-1999. CCAMLR-WG-FSA 99/22. CCAMLR, Hobart.
- Ryan, P.G. and Watkins, B.P. 2002. Reducing incidental mortality of seabirds with an underwater longline setting funnel. *Biological Conservation*, 104: 127–131.
- Sánchez, A., and Belda, E.J. 2003. Bait loss caused by seabirds on longline fisheries in the northwestern Mediterranean: is night setting an effective mitigation measure? *Fisheries Research*, 60: 99–106.
- Seco Pon, J.P., Gandini, P.A. and Favero, M. 2007. Effect of longline configuration on seabird mortality in the Argentine semi-pelagic kingclip *Genypterus blacodes* fishery. *Fisheries Research*, 85: 101–105.
- Smith, N.W.M. 2001. Longline sink rates of an autoline vessel, and notes on seabird interactions. *Science for Conservation* 183. Department of Conservation, Wellington.
- Snell, K.R.S., Brickle, P. and Wolfaardt, A.C. In prep. Quantifying the effectiveness of the Brickle Curtain at preventing foul hooking of seabirds associated with demersal longliners in the Falkland Islands.
- Sullivan, B. 2004. Falkland Islands FAO National Plan of Action for reducing incidental catch of seabirds in longline fisheries. Royal Society for the Protection of Birds.
- Sullivan, B. and Reid, T.A. 2002. Seabird interactions/mortality with longliners and trawlers in Falkland Island waters 2001/02. Falklands Conservation, Stanley, Falkland Islands.
- Weimerskirch, H., Capdeville, D. and Duhamel, G. 2000. Factors affecting the number and mortality of seabirds attending trawlers and long-liners in the Kerguelen area. *Polar Biology*, 23: 236–249.

Other references and resources

- Løkkeborg S. 2008. Review and assessment of mitigation measures to reduce incidental catch of seabirds in longline, trawl and gillnet fisheries. FAO Fisheries and Aquaculture Circular, No. 1040. Rome.
- BirdLife International. 2009. Bycatch mitigation fact-sheets. <http://www.rspb.org.uk/ourwork/policy/marine/international/publications.asp>